

# ABSTRACT

## MATHEMATICAL SCIENCES

HARDEN, KATRINA R. B.A., UNIVERSITY OF ARKANSAS AT LITTLE ROCK,  
1995

### MEAN NUMBER AND VARIANCE OF THE NUMBER OF REAL ZEROS OF A RANDOM TRIGONOMETRIC POLYNOMIAL

Advisor: Dr. J. Ernest Wilkins, Jr.

Thesis dated May, 2002

Suppose that the real numbers  $a_k$  ( $k = 1, 2, \dots, n$ ) are independent random variables, each of which is normally distributed with mean 0 and variance 1. We define the random trigonometric cosine polynomial  $f(x)$  so that

$$f(x) = a_1 \cos x + a_2 \cos 2x + \dots + a_n \cos nx.$$

Let  $N$  be the number of zeros of  $f(x)$  in the real interval  $(0, 2\pi)$ . Then  $N$  is itself a random variable. An explicit formula for the mean value of  $N$  can be found in the literature. An asymptotic expansion of this formula, valid for large values of  $n$ , has also been derived. On the other hand, relatively little is known about the variance of  $N$ . Farahmand has a result that implies a formula for the variance of  $N$  when  $n \geq 4$ . He has used his result to show that the variance  $O(n^{3/2})$  for large  $n$ . This is the best result known at present. We will calculate numerical values for the variance of  $N$  when  $n$  is small, i.e.,  $n = 1, 2$ , and  $3$ .

MEAN NUMBER AND VARIANCE OF THE NUMBER OF REAL ZEROS OF A  
RANDOM TRIGONOMETRIC POLYNOMIAL

A THESIS

SUBMITTED TO THE FACULTY OF CLARK ATLANTA UNIVERSITY  
IN PARTIAL FULFILLMENT OF THE REQUIREMENTS FOR  
THE DEGREE OF MASTER OF SCIENCE

BY

KATRINA RACHELLE HARDEN

DEPARTMENT OF MATHEMATICAL SCIENCES

ATLANTA, GEORGIA

MAY 2002

Riv T19

© 2002

KATRINA R. HARDEN

All Rights Reserved

## ACKNOWLEDGEMENTS

I would first like to give thanks to my Lord and Savior Jesus Christ, who is by my side through all times. I would also like to thank Dr. J. Ernest Wilkins, Jr., who has offered unselfish guidance to me throughout the composition of this thesis. Next, I would like to thank Dr. M. Bleicher for giving me the opportunity to further my education. I would also like to thank Dr. F. Abebe, Dr. H. Adeyemi, and Dr. F. Semwogerere, the readers and faculty members of the Mathematical Sciences Department, for the help and generous amount of time they have given me. Lastly, to Ms. B. Copeland, Mr. D. Torain, and my family, I appreciate all of the love and support you have shown me.

## TABLE OF CONTENTS

ACKNOWLEDGEMENTS.....	ii
LIST OF TABLES.....	iv
Chapter	
1. INTRODUCTION.....	1
2. THE GENERAL CASE.....	3
3. THE CASES WHEN $n = 1$ AND $n = 2$ .....	4
4. THE CASE WHEN $n = 3$ .....	7
5. THE CALCULATION OF $v_3$ AND $\mu_3$ .....	14
WORKS CITED.....	19

## LIST OF TABLES

Table	Page
1. Mean and Variance of $N$ .....	2
2. Regions of Constant $N$ .....	13
3. Java Program for $J_5$ .....	17

## CHAPTER 1

### Introduction

Suppose that  $a_1, a_2, \dots, a_n$  are independent, normally distributed random variables each with mean 0 and variance 1. The number of real zeros of the random trigonometric cosine polynomial,

$$f(x) = a_1 \cos x + a_2 \cos 2x + \dots + a_n \cos nx,$$

in the real interval  $(0, 2\pi)$  is the random variable  $N$ . An explicit formula for the mean value  $\nu_n$  of  $N$  can be found in the literature [2, 3, 8, 9]. An asymptotic expansion of this formula, valid for large values of  $n$ , has also been derived [9]. On the other hand, little is known about the variance  $\sigma_n^2$  of  $N$ . Farahmand [5] has a result when  $n \geq 4$ , and he has deduced an upper bound for  $\sigma_n^2$ . He later [6] obtained a better result that states that  $\sigma_n^2$  is  $O(n^{3/2})$  for large  $n$ . This is the best result known at present. Summaries of much of the information can be found in one or both of the monographs [2, 7].

We will calculate numerical values for the variance of  $N$  when  $n$  is small, i.e.  $n = 1, 2$ , and  $3$ . The calculation is based on an initial explicit determination of  $N$  as a function of the random coefficients, followed by an appropriate integration of  $N$  and  $N^2$  over the probability distribution of the coefficients. This approach is

feasible when  $n$  is small although possibly unsuitable for large values of  $n$ . Integral formulas for the expected values,  $\nu_n$  of  $N$  and  $\mu_n$  of  $N^2$ , when  $n$  is arbitrary, can be found in Chapter 2. Then the variance of  $N$  is  $\sigma_n^2 = \mu_n - \nu_n^2$ . In Chapters 3 and 4 we apply these formulas to the special cases  $n = 1$  and  $n = 2$ , and  $n = 3$ , respectively. After some numerical analysis discussed in Chapter 5, we obtain the results for  $\nu_3$  and  $\sigma_3^2$  that are tabulated in Table 1.

Table 1. MEAN AND VARIANCE OF  $N$

$n$	$\nu_n$	$\sigma_n^2$
1	2	0
2	3	1
3	4.1578	2.4078



## CHAPTER 2

### The General Case

Let  $N(a_1, a_2, \dots, a_n)$  be the number of zeros on the real interval  $(0, 2\pi)$  of the random trigonometric polynomial,

$$f(x) = a_1 \cos x + a_2 \cos 2x + \dots + a_n \cos nx,$$

in which the coefficients  $a_k$  ( $k = 1, \dots, n$ ) are independent real valued random variables, each normally distributed with mean 0 and variance 1, i.e., the probability density function for  $a_k$  is  $(2\pi)^{-\frac{1}{2}} \exp(-\frac{1}{2}a_k^2)$ . Then  $N$  and  $N^2$  are random variables whose respective expected values are the following:

$$\nu_n = (2\pi)^{-n/2} \int_{-\infty}^{\infty} \dots \int_{-\infty}^{\infty} \exp\{-\frac{1}{2}(a_1^2 + a_2^2 + \dots + a_n^2)\} N(a_1, \dots, a_n) da_1 \dots da_n,$$

$$\mu_n = (2\pi)^{-n/2} \int_{-\infty}^{\infty} \dots \int_{-\infty}^{\infty} \exp\{-\frac{1}{2}(a_1^2 + a_2^2 + \dots + a_n^2)\} N^2(a_1, \dots, a_n) da_1 \dots da_n.$$

Moreover, the variance  $\sigma_n^2$  of the random variable  $N$  is  $\mu_n - \nu_n^2$ .

## CHAPTER 3

### The Cases When $n = 1$ and $n = 2$

When  $n = 1$ , the number of real zeros of the random trigonometric polynomial

$$f(x) = a_1 \cos x$$

on the interval  $(0, 2\pi)$  is 2. If  $\nu_1$  is the expected value of  $N(a_1)$ , then  $\nu_1 = 2$ .

Moreover,  $\mu_1 = 4$  and  $\sigma_1^2 = 0$ .

When  $n = 2$ , let  $N(\alpha, \beta; a_1, a_2)$  be the number of real zeros of the random trigonometric polynomial

$$f(x) = a_1 \cos x + a_2 \cos 2x$$

on the interval  $(\alpha, \beta)$ . If  $\nu_2$  is the expected value of  $N(0, 2\pi; a_1, a_2)$ , then

$$\nu_2 = (1/2\pi) \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} N(0, 2\pi; a_1, a_2) \exp\left\{-\frac{1}{2}(a_1^2 + a_2^2)\right\} da_1 da_2.$$

Because  $f(x) = f(2\pi - x)$ , we see that  $N(0, 2\pi; a_1, a_2) = 2N(0, \pi; a_1, a_2)$ . Hence,

$$\nu_2 = (1/\pi) \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} N(0, \pi; a_1, a_2) \exp\left\{-\frac{1}{2}(a_1^2 + a_2^2)\right\} da_1 da_2.$$

Moreover,  $N(0, \pi; a_1, a_2)$  is an even function of both  $a_1$  and  $a_2$ . Therefore,

$$\nu_2 = (4/\pi) \int_0^{\infty} \int_0^{\infty} N(0, \pi; a_1, a_2) \exp\left\{-\frac{1}{2}(a_1^2 + a_2^2)\right\} da_1 da_2.$$

Let  $y = \cos x$ , so that  $\cos 2x = 2y^2 - 1$ . Then

$$f(x) = 2a_2y^2 + a_1y - a_2,$$

so that  $f(x)$  becomes a function  $g(y)$ .

When  $a_1 > 0$  and  $a_2 > 0$ , it is clear that  $g(0) = -a_2 < 0$ ,  $g(1) = a_1 + a_2 > 0$ . Hence, there is a unique  $y_1$  such that  $0 < y_1 < 1$  and  $g(y_1) = 0$ . Because the product of the zeros of  $g(y)$  is  $-1/2$ , the other zero of  $g(y)$  is  $y_2 = -1/(2y_1)$ . This zero is negative because  $y_1$  is positive, and is in the interval  $(-1, 0)$  if and only if  $y_1 > \frac{1}{2}$ . Note that  $g'(y) = 4a_2y + a_1 > 0$  when  $y > 0$ , so that  $y_1 > \frac{1}{2}$  if and only if  $g(\frac{1}{2}) < 0$ . Also note that  $g(\frac{1}{2}) = (a_1 - a_2)/2 < 0$  if and only if  $a_2 > a_1$ . Therefore,  $N(0, \pi; a_1, a_2) = 2$  when  $a_2 > a_1 > 0$ , and  $N(0, \pi; a_1, a_2) = 1$  when  $a_1 > a_2 > 0$ .

Hence,

$$\nu_2 = (8/\pi) \int_0^\infty \left[ \int_{a_1}^\infty e^{-a_2^2/2} da_2 \right] e^{-a_1^2/2} da_1 + (4/\pi) \int_0^\infty \left[ \int_0^{a_1} e^{-a_2^2/2} da_2 \right] e^{-a_1^2/2} da_1.$$

Suppose that  $\rho = \int_{a_1}^\infty e^{-a_2^2/2} da_2$ ,  $w = \int_0^{a_1} e^{-a_2^2/2} da_2$ , so that  $\rho(0) = w(\infty) = \eta = \sqrt{\pi/2}$ . Then

$$\pi\nu_2 = 8 \int_0^\eta \rho \, d\rho + 4 \int_0^\eta w \, dw = 6\eta^2 = 3\pi,$$

so that  $\nu_2 = 3$ . Similarly,

$$\mu_2 = (8/\pi) \int_0^\infty \int_0^\infty N^2(0, \pi; a_1, a_2) \exp\left\{-\frac{1}{2}(a_1^2 + a_2^2)\right\} da_1 da_2,$$

$$\pi\mu_2 = 32 \int_0^\infty \left[ \int_{a_1}^\infty e^{-a_2^2/2} da_2 \right] e^{-a_1^2/2} da_1 + 8 \int_0^\infty \left[ \int_0^{a_1} e^{-a_2^2/2} da_2 \right] e^{-a_1^2/2} da_1$$

$$= 32 \int_0^\eta \rho \, d\rho + 8 \int_0^\eta w \, dw = 20\eta^2 = 10\pi,$$

so that  $\mu_2 = 10$ . Therefore, variance  $\sigma_2^2$  of the random variable N is  $\mu_2 - \nu_2^2 = 1$ .

## CHAPTER 4

### The Case When $n = 3$

When  $n = 3$ , the polynomial to be considered on the interval  $(0, 2\pi)$  is

$$f(x) = a_1 \cos x + a_2 \cos 2x + a_3 \cos 3x.$$

Let  $N(\alpha, \beta; a_1, a_2, a_3)$  be the number of zeros of  $f(x)$  on the real interval  $(\alpha, \beta)$ . If  $\nu_3$  is the expected value of  $N(0, 2\pi; a_1, a_2, a_3)$ , then

$$\nu_3 = (1/2\pi)^{3/2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} N(0, 2\pi; a_1, a_2, a_3) \exp\left\{-\frac{1}{2}(a_1^2 + a_2^2 + a_3^2)\right\} da_1 da_2 da_3.$$

Because  $f(x) = f(2\pi - x)$ , we see that  $N(0, 2\pi; a_1, a_2, a_3) = 2N(0, \pi; a_1, a_2, a_3)$ .

Consequently,

$$\nu_3 = 2^{-1/2} \pi^{-3/2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} N(0, \pi; a_1, a_2, a_3) \exp\left\{-\frac{1}{2}(a_1^2 + a_2^2 + a_3^2)\right\} da_1 da_2 da_3.$$

Let  $y = \cos x$ , so that  $\cos 2x = 2y^2 - 1$  and  $\cos 3x = 4y^3 - 3y$ . Then  $f(x) = 4a_3 G(y; u, v)$ , in which  $G(y; u, v) = y^3 + 2vy^2 + uy - v$ ,  $u = (a_1 - 3a_3)/(4a_3)$ , and  $v = a_2/(4a_3)$ . Let  $N(u, v)$  be the number of real zeros of  $G(y; u, v)$  on the interval  $(-1, 1)$ , so that

$$N(u, v) = N(0, \pi; a_1, a_2, a_3) \text{ if } a_3 \neq 0,$$

$$\nu_3 = 2^{-1/2} \pi^{-3/2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} N(u, v) \exp\left\{-\frac{1}{2}(a_1^2 + a_2^2 + a_3^2)\right\} da_1 da_2 da_3.$$

In order to justify the last equation, we observe that the set of triples  $(a_1, a_2, a_3)$  for which  $a_3 = 0$  is a set of measure zero in the Euclidean space of all triples.

We wish to change the variables of integration from  $a_1, a_2$  to  $u, v$ . For this purpose we need the Jacobian

$$\begin{vmatrix} \partial a_1 / \partial u & \partial a_1 / \partial v \\ \partial a_2 / \partial u & \partial a_2 / \partial v \end{vmatrix}.$$

Because  $a_1 = (4u + 3)a_3$  and  $a_2 = 4a_3v$ , the Jacobian is  $16a_3^2$ . Moreover,  $a_1^2 + a_2^2 + a_3^2 = 2M(u, v)$ , in which  $M(u, v) = 8u^2 + 8v^2 + 12u + 5$ . Therefore,

$$\begin{aligned} \nu_3 &= 2^{-1/2} \pi^{-3/2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} N(u, v) \exp\{-a_3^2 M(u, v)\} 16a_3^2 du dv da_3, \\ \nu_3 &= 2^{7/2} \pi^{-3/2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} N(u, v) \int_{-\infty}^{\infty} \exp\{-a_3^2 M(u, v)\} a_3^2 da_3 du dv. \end{aligned}$$

It is known [1, p. 302, 7.4.4] that  $\int_{-\infty}^{\infty} \omega^2 \exp(-M\omega^2) d\omega = 2^{-1} M^{-3/2} \pi^{1/2}$ . Hence

$$\nu_3 = 2^{5/2} \pi^{-1} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} N(u, v) M(u, v)^{-3/2} du dv.$$

Because  $G(-y; u, -v) = -G(y, u, v)$ , we see that  $N$  is an even function of  $v$ , i.e.,

$N(u, -v) = N(u, v)$ . Therefore,

$$\nu_3 = 2^{7/2} \pi^{-1} \int_{-\infty}^{\infty} du \int_0^{\infty} N(u, v) M(u, v)^{-3/2} dv,$$

and the expected value  $\mu_3$  of  $N^2$  is such that

$$\mu_3 = 2^{9/2}\pi^{-1} \int_{-\infty}^{\infty} du \int_0^{\infty} N^2(u, v) M(u, v)^{-3/2} dv.$$

We observe the following particular values of  $G$ :

$$\begin{aligned} G(-\infty, u, v) &= -\infty, \\ G(-1, u, v) &= v - u - 1, \\ G(0, u, v) &= -v < 0, \\ G(1, u, v) &= v + u + 1, \\ G(+\infty, u, v) &= +\infty. \end{aligned}$$

Descartes' rule of signs [4, p. 77] implies that there exists a unique positive  $y_1$  such that  $G(y_1, u, v) = 0$ . This  $y_1$  lies in the interval  $(0, 1)$  if and only if  $G(1, u, v) = v + u + 1 > 0$ , i.e.,  $u > -v - 1$ . Because  $G(0, u, v) = -v < 0$ , there exists a unique  $y_2$  such that  $-1 < y_2 < 0$  and  $G(y_2, u, v) = 0$  if and only if  $G(-1, u, v) = v - u - 1 > 0$ , i.e.,  $u < v - 1$ . We can now observe that  $N(u, v) = 2$  if  $-v - 1 < u < v - 1$ ; the two zeros are  $y_1$  and  $y_2$ .

If  $u < -v - 1$ , then  $G(1, u, v) < 0$ , which implies that  $y_1 > 1$ . Hence,  $G(y, u, v)$  has no zeros on  $(0, 1)$ . Moreover,  $u < v - 1$  when  $u < -v - 1$ . Hence, there exists a unique  $y_2$  such that  $G(y_2, u, v) = 0$  and  $-1 < y_2 < 0$ . Therefore,  $N(u, v) = 1$  when  $u < -v - 1$ .

Now suppose that  $u > v - 1$  and  $-1 < y < 0$ . Then  $G(y, u, v) < y^3 + 2vy^2 + (v - 1)y - v = (y + 1)[y^2 + (2v - 1)y - v]$ . When  $y = 0$ ,  $y^2 + (2v - 1)y - v = -v < 0$ .

Also when  $y = -1$ ,  $y^2 + (2v - 1)y - v = 2 - 3v$ . We observe that the quadratic polynomial,  $y^2 + (2v - 1)y - v$  is a convex function of  $y$  and its values at the endpoints of the interval  $(-1, 0)$  are respectively the numbers  $2 - 3v$  and  $-v$ , both of which are negative when  $v > 2/3$ . Therefore,  $y^2 + (2v - 1)y - v < 0$  when  $-1 \leq y \leq 0$  and  $v > 2/3$ . Hence,  $G(y, u, v) < 0$  when  $-1 \leq y \leq 0$  and  $v > 2/3$ , and  $G(y, u, v)$  has no zeros on the interval  $(-1, 0)$ . Because  $u > -v - 1$  when  $u > v - 1$ , we know that there is a unique zero  $y_1$  of  $G$  in the interval  $(0, 1)$ . Therefore,  $N(u, v) = 1$  when  $u > v - 1$  and  $v > 2/3$ .

Next, suppose that  $u > v - 1$  and  $0 < v < 2/3$ . We observe that

$$9(\partial G / \partial y) = 9(3y^2 + 4vy + u) = 3(3y + 2v)^2 + 4(3u - 4v^2).$$

Hence,  $(\partial G / \partial y) > 0$  for all  $y$  if  $u > (4v^2)/3$ . Consequently, the maximum of  $G(y, u, v)$  on the interval  $(-1, 0)$  is  $G(0, u, v) = -v < 0$ . Therefore,  $G(y, u, v) < 0$  when  $-1 \leq y \leq 0$  i.e., there are no negative zeros of  $G$  on  $(-1, 0)$ . However, there does exist a unique  $y_1$  such that  $0 < y_1 < 1$ ,  $G(y_1, u, v) = 0$ . Therefore,  $N(u, v) = 1$  when  $u > v - 1$ ,  $0 < v < 2/3$ , and  $u > (4v^2)/3$ . Of course,  $(4v^2)/3 > v - 1$  for all real  $v$ , so that the inequality,  $u > v - 1$ , is superfluous.

Finally, suppose that  $0 < v < 2/3$  and that  $v - 1 < u < (4v^2)/3$ . Then  $\partial G / \partial y = 3y^2 + 4vy + u = 0$  when  $y = y_0$ , in which

$$y_0 = (-2v \pm \sqrt{4v^2 - 3u})/3.$$



Moreover,  $(\partial^2 G)/(\partial y^2) = 6y + 4v = \pm 2\sqrt{4v^2 - 3u}$  when  $y = y_0$ . Therefore,  $G$  has a local maximum when the minus sign is used, so that  $3y_0 = -2v - \sqrt{4v^2 - 3u} < 0$ . Moreover,  $y_0 > -1$  if and only if  $\sqrt{4v^2 - 3u} < 3 - 2v$ . Because  $3 - 2v > 0$  when  $v < 2/3$ , the last inequality holds if and only if  $4v^2 - 3u < (3 - 2v)^2$ , or if and only if  $u > 4v - 3$ . This inequality is surely true because  $u > v - 1 > 4v - 3$  when  $v < 2/3$ . The local maximum value of  $G$  is the maximum value of  $G$  on  $(-1, 0)$ , and is such that

$$27G(y_0, u, v) = 16v^3 - 18uv - 27v + 2(4v^2 - 3u)^{3/2}.$$

Hence,  $G(y_0, u, v) < 0$ , so that  $G(y, u, v) < 0$  when  $-1 \leq y \leq 0$  and  $N(u, v) = 1$ , if and only if

$$2(4v^2 - 3u)^{3/2} < 27v + 18uv - 16v^3.$$

The right hand side of this inequality is greater than  $27v + 18(v - 1)v - 16v^3 = v(9 + 18v - 16v^2)$ , and this is positive because  $0 < v < 2/3$ . Therefore,  $G(y_0, u, v) < 0$  in case

$$4(4v^2 - 3u)^3 < (27v + 18uv - 16v^3)^2,$$

and this inequality holds if and only if,

$$R(u, v) = u^3 - u^2v^2 + 9uv^2 + v^2(6.75 - 8v^2) > 0.$$

We see that  $3(\partial R/\partial u) = 3(3u^2 - 2v^2u + 9v^2) = (3u - v^2)^2 + v^2(27 - v^2) > 0$  when  $v^2 < 27$ . Moreover,  $R(0, v) = v^2(6.75 - 8v^2) > 0$  when  $0 < v < 2/3$ , and  $R(v-1, v) = -(3v-2)^2(4v^2+1) < 0$  when  $v \neq 2/3$ . Therefore, when  $0 < v < 2/3$ , there exists a unique real number  $F(v)$  such that  $R(F(v), v) = 0, v-1 < F(v) < 0$ . Hence,  $R(u, v) > 0$  when  $u > F(v)$ , and  $R(u, v) < 0$  when  $u < F(v)$ . Accordingly,  $G(y_0, u, v) < 0$  when  $F(v) < u < (4v^2)/3$  and  $G(y_0, u, v) > 0$  when  $v-1 < u < F(v)$ . Therefore,  $N(u, v) = 1$  when  $F(v) < u < (4v^2)/3$  and  $0 < v < 2/3$ .

When  $v-1 < u < F(v)$ , however,  $G(y_0, u, v) > 0$ ,  $G(-1, u, v) < 0$ , and  $G(0, u, v) < 0$ . Hence, there are exactly two zeros of  $G$  on the interval  $(-1, 0)$ , and so  $N(u, v) = 3$  when  $v-1 < u < F(v)$  and  $0 < v < 2/3$ .

In order to find  $F(v)$ , we suppose that  $v = c^{3/2}$ ,  $F = cz$ . Then  $z^3 + pz^2 + qz + r = 0$ , in which  $p = -c^2$ ,  $q = 9c$ , and  $r = 6.75 - 8c^3$ . Using Cardan's formulas [4, p. 42-44], we see that  $z = A + B - (p/3)$ , in which  $a = (3q - p^2)/9 = c(27 - v^2)/9$ ,  $b = (9pq - 2p^3 - 27r)/54 = (8v^4 + 540v^2 - 729)/216$ ,  $B^3 = b - (b^2 + a^3)^{1/2} = b - \{(8v^2 + 27)/12\}^{3/2}$ ,  $A = -a/B$ . The value of  $F(v)$  can be calculated, for a given value of  $v$ , from these formulas.

We summarize our calculation of  $N(u, v)$  in Table 2.

Table 2. REGIONS OF CONSTANT  $N$ 

$$N(u, v) = 1 \text{ when } 0 < v < \infty \text{ and } u < -v - 1,$$

$$\text{when } 0 < v < 2/3 \text{ and } u > F(v),$$

$$\text{and when } v > 2/3 \text{ and } u > v - 1;$$

$$N(u, v) = 2 \text{ when } 0 < v < \infty \text{ and } -v - 1 < u < v - 1;$$

$$N(u, v) = 3 \text{ when } 0 < v < 2/3 \text{ and } v - 1 < u < F(v).$$

## CHAPTER 5

### The Calculation of $\nu_3$ and $\mu_3$

To calculate  $\nu_3$  and  $\mu_3$  we will begin with the indefinite integral,

$$\begin{aligned} K(u, v) &= \int [M(u, v)]^{-3/2} du = \int [8u^2 + 8v^2 + 12u + 5]^{-3/2} du \\ &= (4u + 3)/[2(16v^2 + 1)(8u^2 + 8v^2 + 12u + 5)^{1/2}]. \end{aligned}$$

With the help of Table 2, we now see that

$$\begin{aligned} 2^{-7/2}\pi\nu_3 &= \int_0^\infty [K(-v-1, v) - K(-\infty, v)]dv + \int_0^{2/3} [K(\infty, v) - K(F(v), v)]dv \\ &\quad + \int_{2/3}^\infty [K(\infty, v) - K(v-1, v)]dv + 2 \int_0^\infty [K(v-1, v) - K(-v-1, v)]dv \\ &\quad + 3 \int_0^{2/3} [K(F(v), v) - K(v-1, v)]dv. \end{aligned}$$

Therefore,  $2^{-7/2}\pi\nu_3 = J_1 + J_2 + J_3 + J_4 + J_5$ , in which

$$J_1 = \int_0^\infty K(\infty, v) dv - \int_0^\infty K(-\infty, v) dv,$$

$$J_2 = - \int_0^\infty K(-v-1, v) dv,$$

$$J_3 = \int_0^\infty K(v-1, v)dv,$$

$$J_4 = -2 \int_0^{2/3} K(v-1, v)dv,$$

$$J_5 = 2 \int_0^{2/3} K(F(v), v)dv.$$

Note that  $K(\infty, v) = -K(-\infty, v) = 2^{-1/2}(16v^2+1)^{-1}$ , and  $\int_0^\infty (16v^2+1)^{-1}dv = \pi/8$ . Hence  $J_1 = (\sqrt{2}\pi)/8$ .

If we make the sequence of changes of variables of integration,

$$4v = \xi, \quad \xi + 1/2 = (\sqrt{3}/2) \tan \theta, \quad \sin (\theta - \pi/3) = t/\sqrt{2},$$

we find that

$$\begin{aligned} J_2 &= \int_0^\infty \frac{(4v+1) dv}{2(16v^2+1)(16v^2+4v+1)^{\frac{1}{2}}} \\ &= 2^{-\frac{5}{2}} \int_{-2^{-1/2}}^{2^{-1/2}} (1+t^2)^{-1} dt = 2^{-\frac{3}{2}} \tan^{-1}(2^{-1/2}) \approx 0.2176049379. \end{aligned}$$

Similarly, if

$$4v = \xi, \quad \xi - 1/2 = (\sqrt{3}/2) \tan \theta, \quad \sin (\theta + \pi/3) = t/\sqrt{2},$$

we find that

$$\begin{aligned} J_3 &= -2^{-5/2} \int_{2^{-1/2}}^{2^{-1/2}} (1+t^2)^{-1} dt = 0, \\ J_4 &= 2^{-\frac{3}{2}} \int_{-2^{-1/2}}^s (1+t^2)^{-1} dt = 2^{-\frac{3}{2}} [\tan^{-1}(s) - \tan^{-1}(2^{-1/2})] \approx 0.513885010, \end{aligned}$$

in which  $s = (11\sqrt{2})/14$ . Note that  $\tan^{-1}(s) \approx 0.838006592$ .

Unfortunately, it does not appear that the integral in  $J_5$  is elementary. Nonetheless, we can use Simpson's rule with sufficiently small interval size to get an accurate approximation to  $J_5$ . This approximation was computed by the JAVA computer program listed in table 3. This yields the result that  $J_5 \approx 0.3028975974$ . We can now use the numerical values of  $J_1$ ,  $J_2$ ,  $J_3$ ,  $J_4$ , and  $J_5$  to find that

$$\nu_3 = 4.15779773.$$

Moreover,

$$\begin{aligned}
2^{-7/2}\pi\mu_3 &= \int_0^\infty [K(-v-1, v) - K(-\infty, v)] dv + \int_0^{2/3} [K(\infty, v) - K(F(v), v)] dv \\
&\quad + \int_{2/3}^\infty [K(\infty, v) - K(v-1, v)] dv + 4 \int_0^\infty [K(v-1, v) - K(-v-1, v)] dv \\
&\quad + 9 \int_0^{2/3} [K(F(v), v) - K(v-1, v)] dv \\
&= J_1 + 3J_2 + 3J_3 + 4J_4 + 4J_5,
\end{aligned}$$

so that

$$\mu_3 = 4 - \frac{8}{\pi} \tan^{-1}(2^{-1/2}) + \frac{32}{\pi} \tan^{-1} s + \frac{2^{13/2}}{\pi} J_5 \approx 19.69507562.$$

Consequently,

$$\sigma_3^2 = \mu_3 - \nu_3^2 \approx 2.40779366,$$

and

$$\sigma_3 = 1.551706694.$$

Table 3. JAVA PROGRAM FOR  $J_5$ 

```

public class ComputationOfJ5 {
    public static void main (String [] args) {
        double q, v, sum, integralSolution, pv;
        double h, prevIntegralSol, difference;

        h = 1/12D;

        prevIntegralSol = 0;

        integralSolution = 0;

        difference = 0;

        do {

            prevIntegralSol = integralSolution;

            q = 4.0;

            v = h;

            sum = 3/(Math.sqrt(5.0));

            do {

                pv = calculatepv(v);

                sum = sum + q * pv;

                v = v+h;

                q = 6-q;

            } while (v < (2/3D) + (h/2D));

```

Table 3 - Continued

```

sum = sum - pv;

integralSolution = h/3 * sum;

h = h/8;

difference = Math.abs (integralSolution - prevIntegralSol);

} while (difference > 1/1000000000000D );

System.out.println ("integral solution = " + integralSolution);

}

public static double calculatepv (double v) {

double u = Math.pow (v, 2);

double c = Math.pow (u, 1/3D);

double t = Math.pow (((8*u + 27) / 12), 3/2D);

double d = (c * (27 - u) ) / 9;

double w = (729 - u * (8*u + 540) ) / 216;

double b = - (Math.pow ((w + t), 1/3D) );

double a = - (d / b);

double f = (a + b) * c + (u / 3);

double pv = ((4*f) + 3) / ((16*u+1) * Math.sqrt (f*(8*f + 12) + 5+8*u));

return pv;

}

}

```



## WORKS CITED

1. Abramowitz, M. and I. A. Stegun, eds. *Handbook of Mathematical Functions and Tables*. New York: Dover Publications, 1965.
2. Bharucha-Reid, A. T. and M. Sambandham. *Random Polynomials*. Orlando: Academic Press, 1986.
3. Das, M. The average number of real zeros of a random trigonometric polynomial. *Proceedings of the Cambridge Philosophical Society* 64 (1968): 721-729.
4. Dickson, L. E. *New First Course in the Theory of Equations*. New York: Wiley Publishing Company, 1939.
5. Farahmand, K. On the variance of the number of real roots of a random trigonometric polynomial. *Journal of Applied Mathematics and Stochastic Analysis* 3 (1990): 253-262.
6. Farahmand, K. On the variance of the number of real zeros of a random trigonometric polynomial. *Journal of Applied Mathematics and Stochastic Analysis* 10 (1997): 57-66.
7. Farahmand, K. *Topics in Random Polynomials*. Pitman Research Notes in Mathematics Series 393. Edinburgh Gate: Addison Wesley Longman Ltd, 1998.
8. Sambandham, M. On the number of real zeros of a random trigonometric polynomial. *Transactions of the American Mathematical Society* 238 (1978): 57-70.
9. Wilkins, J. E. Mean number of real zeros of a random trigonometric polynomial. *Proceedings of the American Mathematical Society* 111 (1991): 851-863.